

Preventive health guidelines

2024

Take steps today for a healthier future

Wellpoint Washington, Inc. pays for certain tests to find diseases early, routine wellness exams, and shots to help you and your family stay well. This is called preventive care.

These guidelines are based on state-specific requirements and tips from health experts, including:

- American Academy of Family Physicians (AAFP)
- American Academy of Pediatrics (AAP)
Bright Futures
- Advisory Committee on Immunization Practices (ACIP)
- American College of Obstetricians and Gynecologists (ACOG)
- American Cancer Society (ACS)
- Centers for Disease Control and Prevention (CDC)
- U.S. Preventive Services Task Force (USPSTF)



Your plan may not pay for all the services and treatments listed.

To learn more about what Wellpoint covers, either:

- Check the member handbook.
- Call Member Services at the number on your member ID card.
- Visit wellpoint.com/wa/medicaid.

Always get personal medical advice from your doctor.

This guide does not mention every condition and treatment. Ask the doctor which exams, tests, and vaccines are right for you or your child, when to receive them, and how often.

Well-baby visits — birth to 2 years old

Infants need to be seen by a doctor at birth, at the following ages, and as the doctor suggests:

- 3–5 days old
- 1 month
- 2 months
- 4 months
- 6 months
- 9 months
- 12 months
- 15 months
- 18 months
- 24 months

Babies who leave the hospital less than two days (48 hours) after birth need to be seen by a doctor within 2 to 4 days after being born.

A well-baby visit may include the following:

- A full-body exam
- Vaccines
- Other tests and screenings as needed, listed below
- Talking about:
 - Newborn care, safety, and development
 - Nutrition and feeding
 - Parent and family health and well-being

Screenings	When to receive them
Weight, length, and head measurement	At each visit
BMI percentile*	At 24 months
Newborn metabolic, such as PKU (when the body is unable to break down protein), sickle cell (an inherited blood disorder), and thyroid screening	Birth to 2 months old (best checked at 3 to 5 days old). Bilirubin at birth (checks for liver problems).
Critical congenital heart defect (birth defects of the heart)	At 0 to 6 weeks old
Growth and development	At 9 to 11 months, 18 months, and 30 months. Monitor at 0 to 6 weeks, 2 to 3 months, 4 to 5 months, 6 to 8 months, 12 to 14 months, and 15 to 17 months.
Behavioral and mental health	At each visit
Hearing	At 0 to 6 weeks. Check for risks at all other visits and follow up if needed.
Vision	At 2 and 24 months. Use instrument-based screenings to check for risks.
Blood pressure	Check for risks at each visit
Oral and dental health	Check oral health and refer to a dentist, if needed — begin yearly dental exams starting at 12 months. Fluoride varnish when teeth start coming in (usually around 6 to 24 months old). Fluoride prescription based on your drinking water (from 6 to 24 months old).
Hemoglobin or hematocrit (blood count)	Once between 9 to 11 months or 12 to 24 months. Check for risks at all other visits.
Lead testing	At 12 and 24 months old. Check for risks as the doctor suggests.
Lipid disorder (cholesterol problems)	Check for risks at 24 months

Well-baby visits — birth to 2 years old *continued*

Screenings	When to receive them
Autism (a condition that affects social skills and the way one communicates)	At 18 and 24 months
Maternal postpartum depression (after a mother gives birth)	At 6 months
Tuberculosis	At 0 to 6 weeks, 6 to 8 months, 12 to 14 months, and 24 months

* Height and weight are used to check body mass index (BMI). Checking someone's BMI helps determine if they are a healthy weight for their height, or if they are under or over weight.

Well-child visits — 2 1/2 to 10 years old

Depending on your child's age, the doctor may talk with you about:

- Health history.
- How to promote healthy nutrition.
- Exercise, growth, safety, and healthy habits.
- Any learning or school issues.
- Emotional and mental health.
- Family and home living issues.

During the annual visit, your child may receive:

- A full-body exam.
- Vaccines.
- Other tests and screenings.

Screenings	When to receive them
Height, weight, BMI percentile*	At each visit
Development — brain, body, and behavior	At each visit
Vision	At each visit
Hearing	At each visit
Oral and dental health	Referral to a dentist, if needed. Check oral health and have a dental exam each year. Fluoride varnish, fluoride treatment, and checking for risks at each visit (between 2 to 10 years old). Follow up if needed.
Lead testing	Check for risks each year through age 6
Tuberculosis	Check for risks and test as the doctor suggests
Hemoglobin or hematocrit (blood count)	Check for risks each visit
Blood pressure	Each year
Lipid disorder (cholesterol problems)	Once between ages 9 to 10 Check for risks at each visit

* Height and weight are used to check body mass index (BMI). Checking someone's BMI helps determine if they are a healthy weight for their height, or if they are under or over weight.

Well-child visits — 11 to 20 years old

Depending on age, the doctor may talk about:

- **Health history.**
- **Growth and development**, such as oral health habits, body image, healthy eating, physical activity, and sleep.
- **Emotional well-being**, including mood control and overall mental health.
- **Safe sex**, especially reducing risks of sexually transmitted infections and diseases (STIs and STDs) and pregnancy.
- **Substance use**, whether that be drinking alcohol or using tobacco, e-cigarettes, or prescription or illegal drugs.
- **School performance.**
- **Family and home living issues.**
- **Safety**, such as seat belt use, helmet use, and sun protection.
- **Firearm safety**, if you own or are around guns.

During the annual visit, the doctor may give:

- A full-body exam.
- Vaccines.
- Other tests and screenings.

Screenings	When to receive them
Height, weight, BMI*	Percentile to age 18, then BMI each year
Development — mind, body, and behavior	At each visit
Depression	At each visit
Blood pressure	At each visit
Vision	At ages 11 to 12, and 15 to 16 years old Check for risks from ages 13 to 14, 17 to 18, and 19 to 20
Hearing	Once between 11 to 14, 15 to 16, and 19 to 20 years old Check for risks from ages 17 to 18
Oral and dental health	Each year Check for risks of needed fluoride prescription between ages 11 to 12, 13 to 14, and 15 to 16
Tuberculosis	Check for risks each year
Hemoglobin or hematocrit (blood count)	Check for risks each year
Lipid disorder (cholesterol problems)	Once between ages 9 to 11 Once between ages 17 to 18 Check for risks at all other visits
STIs, including chlamydia	Starting at age 11, if sexually active, then check for risks at each visit
HIV	Screen once between ages 15 to 18 Check for risks at each visit
Substance use disorder and tobacco addiction	Check for risks each year

* Height and weight are used to check body mass index (BMI). Checking someone's BMI helps determine if they are a healthy weight for their height, or if they are under or over weight.

Wellness visits — adult women

During your visit, the doctor may talk with you about:

- Health history.
- Diet and physical activity.
- Mental health, including depression.
- Oral and dental health.
- Tobacco use, or how to quit.
- Avoiding secondhand smoke.
- Drinking alcohol or substance use.
- Skin cancer risks.
- Family planning, including:
 - Safe sex.
 - Birth control to help avoid unwanted pregnancy.
 - Spacing out pregnancies to have the best birth outcomes.
 - Checking for sexually transmitted infections and diseases (STIs and STDs), including HIV and hepatitis B (if at risk).
 - Folic acid supplements for women of childbearing age.

You may also receive vaccines and these screenings:

Screenings	When to receive them
Height, weight, BMI*	Each year or as your doctor suggests
Blood pressure	Each year or as your doctor suggests. Recheck high readings at home.
Mammogram** (breast X-ray)	Each year for ages 40 to 65+ Consider screening every 2 years from ages 50 to 74
Cervical cancer	For ages 21 to 29, Pap test every 3 years For ages 30 to 65, either do a Pap test every 3 years or an HPV test alone, or a combination Pap test and HPV test every 5 years Stop testing at age 65 if the last 3 Pap tests or last 2 co-tests (Pap plus HPV) within the last 10 years were normal. If there was an abnormal Pap test within the past 20 years, talk with your doctor. From ages 50 to 75***, your doctor may suggest one or more of these test options:
Colorectal cancer (of the colon and rectum)	Stool (feces) tests: <ul style="list-style-type: none"> • Fecal immunochemical test (FIT) • FIT-DNA: stool and DNA combo test • Guaiac-based fecal occult blood test (gFOBT) Visual tests: <ul style="list-style-type: none"> • Colonoscopy (using a small camera on the end of a flexible tube to look at your entire colon) • CT colonography (using a CT scanner to take images of inside the colon) • Flexible sigmoidoscopy (using a small camera on the end of a flexible tube to look at the last part of your colon, called the sigmoid colon)
Chlamydia and gonorrhea	If sexually active and age 24 or younger

Wellness visits — adult women *continued*

Screenings	When to receive them
Cholesterol	Statins (cholesterol medicine) may be needed for people ages 40 to 75 who have a higher risk of cardiovascular disease (such as heart disease)
Glucose (blood sugar) screening for type 2 diabetes	As your doctor suggests from ages 40 to 70, especially if overweight or obese. Individuals with high blood sugar should talk to their doctor about intensive counseling interventions to promote a healthy diet and physical activity.
Hepatitis C	Screen between the ages of 18 and 79 years Testing should start no later than age 65
Osteoporosis (checks how dense your bones are)	Women in menopause should talk to their doctor about osteoporosis and have the test if at risk

* Height and weight are used to check body mass index (BMI). Checking someone's BMI helps determine if they are a healthy weight for their height, or if they are under or over weight.

** Women should talk to their doctor and make a personal choice about the best age to start having mammograms and possibly screen every two years when older.

*** If you are African-American, screening can begin at age 45.

Pregnant individuals

Within the first three months of pregnancy, it's important to visit a doctor to set up a prenatal care plan. At each visit, your doctor will check your health and the health of your baby. The doctor may talk to you about:

- Health history.
- What to eat.
- How to be active when pregnant.
- Avoiding tobacco, drugs, alcohol and other substances.
- Breastfeeding, lactation supplies and counseling.

Testing:

Based on your past health, your doctor may want you to have these screenings:

- **Depression** screenings (done during and after pregnancy)
- **Diabetes**
- **Preeclampsia*** (high blood pressure that causes other problems during pregnancy)
- **Hematocrit/hemoglobin** (blood count)
- **Rubella immunity** (to find out which women need the rubella, aka German measles, vaccine after giving birth)
- **Rh(D) blood type and antibody testing** (checks to see if your blood type and your baby's blood type are compatible.) If Rh(D) negative, repeat test at 24 to 28 weeks.
- **Hepatitis B**
- **HIV**
- **Syphilis**
- **Urine** for asymptomatic bacteriuria, as your doctor suggests

Other tests and screenings:

- **Amniocentesis** (an ultrasound and testing of the fluid surrounding your baby)
- **Cell-free DNA** (a blood test to check for chromosomal abnormalities in the baby)
- **Chorionic villus sampling** (checks for birth defects and more)
- **Ultrasound tests** (to look at the baby in the womb.) During the first three months, these are done along with blood tests to check the baby for chromosomal abnormality risk and more.

These and other tests can check the baby for health concerns. The right tests and the right times to do them depend on:

- Your age.
- Your medical history and family history.

Talk to your doctor about:

- Which tests may be best for you.
- What the tests can tell you about your baby.
- Any risks.

* If you have a high risk of preeclampsia, your doctor may recommend taking a low-dose aspirin to prevent other problems while you are pregnant.

Vaccines:

- **Flu:** If you are pregnant during flu season (October through March), your doctor may want you to have the inactivated (killed) flu shot.
- **Tdap:** Pregnant teens and adults need a Tdap vaccine during each pregnancy. It's best to receive the vaccine between weeks 27 and 36, although it may be given at any time during pregnancy.

It's best to receive most vaccines before pregnancy. Women should check with their doctor to make sure their vaccines are up to date.

You should NOT receive these vaccines while you are pregnant:

- **Measles, mumps, rubella (MMR)**
- **Varicella (chickenpox)**

Wellness visits — adult men

During your visit, the doctor may talk with you about:

- Health history.
- Diet and physical activity.
- Mental health, including depression.
- Oral and dental health.
- Tobacco use, or how to quit.
- Avoiding secondhand smoke.
- Drinking alcohol and substance use.
- Skin cancer risks.
- Family planning, including:
 - Safe sex and preventing unwanted pregnancy with a partner.
 - Checking for sexually transmitted infections and diseases (STIs and STDs), including HIV and hepatitis B (if high risk).

You may also receive vaccines and these screenings:

Screenings	When to receive them
Height, weight, BMI*	Each year or as your doctor suggests
Abdominal aortic aneurysm (enlarged blood vessels in the abdomen)	Once between ages 65 to 75 if you have ever smoked
Blood pressure	Each year or as your doctor suggests. Recheck high readings at home.
Cholesterol	Statins (cholesterol medicine) may be needed for people ages 40 to 75 who have a higher risk of cardiovascular disease (such as heart disease) From ages 50 to 75**, your doctor may suggest one or more of these test options: Stool (feces) tests: <ul style="list-style-type: none">• Fecal immunochemical test (FIT)• FIT-DNA: stool and DNA combo test• Guaiac-based fecal occult blood test (gFOBT) Visual tests: <ul style="list-style-type: none">• Colonoscopy (using a small camera on the end of a flexible tube to look at your entire colon)• CT colonography (using a CT scanner to take images of inside the colon)• Flexible sigmoidoscopy (using a small camera on the end of a flexible tube to look at the last part of your colon, called the sigmoid colon)
Colorectal cancer (of the colon and rectum)	
Glucose (blood sugar) screening for type 2 diabetes	As your doctor suggests from ages 40 to 70, especially if overweight or obese. Individuals with high blood sugar should talk to their doctor about intensive counseling interventions to promote a healthy diet and physical activity.
Hepatitis C	Screen between the ages of 18 and 79 years
Prostate cancer	From ages 55 to 69, talk with your doctor about the risks and benefits of prostate cancer tests

* Height and weight are used to check body mass index (BMI). Checking someone's BMI helps determine if they are a healthy weight for their height, or if they are under or over weight.

** If you are Black, screening can begin at age 45.

Suggested vaccine schedule

For more information about vaccines, visit [cdc.gov/vaccines](https://www.cdc.gov/vaccines). Coronavirus disease 2019 (COVID-19) vaccines as recommended by the CDC ([cdc.gov/vaccines/hcp/ac](https://www.cdc.gov/vaccines/hcp/ac))

Vaccines ↓ Ages →	Birth	1-2 months	2 months	4 months	6 months	6-18 months	12-15 months	15-18 months	19-23 months	4-6 years	11-12 years	13-18 years	19-64 years	65+ years
Hepatitis B	✓	✓				✓								
Rotavirus (RV)			2-dose or 3-dose series											
Diphtheria, tetanus, pertussis (DTaP)			✓	✓	✓			✓		✓				
Tetanus, diphtheria, pertussis (Td/Tdap)											Tdap		Every 10 years	
Haemophilus influenzae type b (Hib)			3-4 doses between 2 to 15 months with 1st dose at 2 months, last dose at 12 to 15 months											
Pneumococcal conjugate (PCV)			✓	✓	✓		✓							
Inactivated polio virus (IPV)			✓	✓		✓				✓				
Influenza (flu)					Suggested each year from 6 months to 65+ years; 2 doses at least 4 weeks apart are recommended for children between 6 months to 8 years old having the vaccine for the first time									
Measles, mumps, rubella (MMR)							✓			✓				
Varicella (chickenpox)							✓			✓				
Hepatitis A							2-dose series between 12 to 23 months; taken 6 to 18 months apart							
Human papillomavirus (HPV)											2-dose series			
Meningococcal											✓	Booster at age 16; MenB-FHb at ages 16 to 23		
Pneumococcal 13-valent conjugate (PCV13)														✓
Pneumococcal polysaccharide (PPSV23)														✓
Zoster (HZ/su) recombinant vaccine														2-dose series for ages 50+; 2 to 6 months apart

* For more information about updated HPV vaccines, see the Centers for Disease Control and Prevention website: *Use of a 2-Dose Schedule for Human Papillomavirus Vaccination — Updated Recommendations of the Advisory Committee on Immunization Practices* (December 16, 2016): [cdc.gov/mmwr/volumes/65/wr/mm6549a5.htm](https://www.cdc.gov/mmwr/volumes/65/wr/mm6549a5.htm).

Hepatitis A (ages 2 to 18): If you or your child has not had this vaccine before, talk to your doctor about a catch-up vaccine.

Hepatitis B: The first dose should be given within 24 hours of birth if the birth was outside of a hospital. Children may receive an extra dose (four-dose series) at 4 months if the combination vaccine is used after the birth dose.

Rotavirus (RV): Receive a two-dose or three-dose series (depending on the brand of vaccine used).

Tdap (children through adults): If you or your child (age 7 or older) never received this vaccine, talk to the doctor about a catch-up vaccine.

Haemophilus influenzae type b (Hib): Receive a three-dose or four-dose series (depending on the brand of vaccine used).

Pneumococcal conjugate (PCV): Children ages 14 months to 59 months who received an incomplete PCV13 series.

Influenza (flu): Visit [flu.gov](https://www.flu.gov) or [cdc.gov](https://www.cdc.gov) to learn more about this vaccine. Children 6 months to 8 years having the vaccine for the first time should have two doses four weeks apart.

Measles, mumps, rubella (MMR) and varicella (chickenpox): Teens and adults should be up to date on their MMR vaccines. Chickenpox vaccines are for children who have not had chickenpox.

Human papillomavirus (HPV): * Children who are 11 to 12 years old receive two doses of the HPV vaccine at least six months apart. (The vaccine series can start at age 9.) Teens and young adults who start the series later (at ages 15 to 26) need three doses of HPV vaccine to protect against cancer-causing HPV infection. Adults ages 27 to 45 should talk to their doctor to see if an HPV vaccine is right for them.

Meningococcal: When given to healthy teens who are not high risk for meningococcal disease, two doses of MenB-FHbp should be given six months apart. This timing is very important. If a second dose is given before six months, a third dose should be given six months after the first dose. For persons at high risk for meningococcal disease and during serogroup B outbreaks, three doses of MenB-FHbp should be given. The second and third dose should be given 1 to 2 months and six months after the first dose.

Pneumococcal 13-valent conjugate (PCV13)/Pneumococcal polysaccharide (PPSV23): Adults age 65 and older and certain adults younger than 65 who are at risk should receive both a PCV13 and PPSV23. Ask your doctor what dose is best for you.

Zoster: Two doses of the Shingrix (HZ/su) vaccine, given 2 to 6 months apart, is recommended for adults 50 and older, including those who received the Zostavax (shingles) vaccine.



Do you need help with your healthcare, talking with us, or reading what we send you? We provide our materials in other languages and formats at no cost to you. Call us toll free at 833-731-2167 (TTY 711).

¿Necesita ayuda con su cuidado de la salud, para hablar con nosotros o leer lo que le enviamos? Ofrecemos nuestros materiales en otros idiomas y formatos sin costo alguno. Llámenos a la línea gratuita 833-731-2167 (TTY 711).

Wellpoint Washington, Inc. complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability or sex.