

Screening Brief Intervention Referral to Treatment (SBIRT)

Medicaid Network Growth and Strategy

What is SBIRT?

Screening (S)

A very brief set of questions that identifies risk of substance use disorder (SUD)-related problems:

- Should last 5-10 minutes.
- Reimbursement requires use of validated screening instruments.

Brief Intervention (BI)

A short (5-20 minutes) counseling session that raises awareness of risks and motivates the client toward acknowledgement of the problem:

- Uses motivational interviewing techniques to encourage lifestyle change.

Referral to Treatment (RT):

- Warm hand-off to a provider who can provide specialized treatment to the patient.



Potential benefits for patients

Positively affects

- Patients with substance use disorders (SUDs)
- Patient morbidity and mortality rates

Reduces

- Healthcare costs
- Work impairment and incidents of driving under the influence

Improves

- Access to treatment
- Neonatal and post-partum outcomes



Source: Substance Use and Mental Health Services Administration (SAMHSA). Screening, Brief Intervention, and Referral to Treatment (SBIRT). 2023. <https://www.samhsa.gov/sbirt>

Potential benefits for providers

Awareness

- Increases clinicians' awareness of substance use issues

Better approach

- Offers clinicians a more systematic approach to addressing substance use, identifying more *hidden* cases

Cost-effectiveness

- Studies have shown that for every \$1 spent, SBIRT for alcohol use saves \$2-\$4



Who can provide SBIRT?



Most effective in:


- Primary care centers
- Emergency rooms (ER) and trauma centers
- Community health settings

Healthcare workers who can provide SBIRT:


- Primary care providers (MD/DOs, PAs, ARNPs)
- Behavioral health providers (therapists, counselors, psychiatrists, clinical social workers)
- OB/GYNs and midwives
- Pediatricians
- Nurses
- Any provider in nearly any setting!



Example ratios



Decision Tree (example)





Does SBIRT work?

Yes! SBIRT is an evidence-based practice.

Does SBIRT work?

Yes! SBIRT is an evidence-based practice.

Project TrEAT: Trial of Early Alcohol Treatment

The program included 17 primary care practices comprised of 64 physicians.

Approximately 18,000 patients were screened:

- Around 500 men and 300 women screened positive for at-risk drinking.
- They were randomized into two groups of approximately 400 each and followed for 48 months.

Both the control and intervention group received a general health booklet with information about seat belt use, immunizations, exercise, tobacco, alcohol, and drugs.

The intervention group also received two 10-15-minute sessions by a primary care physician (PCP) using a scripted workbook.



Project TrEAT Statistics

Utilization (post-intervention)	SBIRT	Control
ER visits	302	376
Days of hospitalization	420	664
Patients considered heavy drinkers		
Baseline	46.7%	49.2%
12 months post-intervention	20.1%	33.5%
Patients reporting binge drinking		
Baseline	85.0%	86.9%
36 months post-intervention	57.4%	71.5%





SBIRT components

Prescreening

Prescreening is a very quick approach to identifying people who need a longer screen or brief intervention or treatment.

Self-report:

- Patient discloses concern about their alcohol or drug use.

Provider questions:

- *How many times in the past month have you had X or more drinks in a day?*
- *How many times in the past month have you used an illegal drug or used a prescription medication for nonmedical reasons?*

Biological:

- Blood alcohol level test
- Urine screening for drugs



How is risk defined?

At-risk alcohol use is defined as:

Drinks	Men	Women	65+
Per occasion	> 4	> 3	> 1
Per week	> 14	> 7	> 7

Any illicit substance use reported should be followed by a full screening.



Screening tools guidelines

Characteristics of a good screening tool:

- Brief (10 or fewer questions)
- Flexible
- Easy to administer and easy for the patient
- Addresses alcohol and other drug use
- Indicates need for further assessment or intervention
- Has good sensitivity and specificity



Screening tools (cont.)

Screening tool ⁵	Age range or population	Overview
Alcohol Use Disorder Identification Test (AUDIT) ¹	All patients	Developed by the World Health Organization (WHO). Appropriate for all ages, genders, and cultures.
Alcohol, Smoking, and Substance Abuse Involvement Screen Test (ASSIST) ²	Adults	Developed by the WHO. Simple screener for hazardous use of substances (including alcohol, tobacco, other drugs).
Drug Abuse Screening Test (DAST-10) ³	Adults	Screener for drug involvement, does not include alcohol, during last 12 months.
Car, Relax, Alone, Forget, Family or Friends, Trouble (CRAFT) ⁴	Adolescents	Alcohol and drug screening tool for patients under 21. Recommended by American Academy of Pediatrics.
Bold indicates our recommended screening tools.		

Babor, T. F., & Grant, M. (1989). From clinical research to secondary prevention: international collaboration in the development of the Alcohol Disorders Identification Test (AUDIT). *Alcohol Health & Research World*, 13(4), 371+.

Group, W.A.W. (2002), The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): development, reliability and feasibility. *Addiction*, 97: 1183-1194.

Skinner, Harvey A. (2002), The drug abuse screening test. *Addictive Behaviors*, 7(4): 363-371.

Knight JR, Shrier LA, Bravender TD, Farrell M, Vander Bilt J, Shaffer HJ. A new brief screen for adolescent substance abuse. *Arch Pediatr Adolesc Med*. 1999 Jun;153(6):591-6



Screening tools (cont.)

Screening tool	Age range or population	Overview
Screening to Brief Intervention (S2BI) ¹	Adolescents	Assesses frequency of alcohol and substance use, for patients ages 12-17.
NIAAA Alcohol Screening for Youth ²	Adolescents and children	Two-item scale to assess alcohol use (self and friends/family), for patients ages 9-18.
Tolerance, Worried, Eye Opener, Amnesia, K/Cut Down (TWEAK) ³	Pregnant women	Five-item scale to screen for risky drinking during pregnancy. Recommended for OB/GYNs.
Substance Use Risk Profile-Pregnancy (SURP-P) ⁴	Pregnant women	Three-item scale to screen for drug use during pregnancy. Recommended for OB/GYNs.
Bold indicates our recommended screening tools.		

Levy, S., Weiss, R., Sherritt, L., Ziemnik, R., Spalding, A., Van Hook, S., & Shrier, L. A. (2014). An electronic screen for triaging adolescent substance use by risk levels. JAMA Pediatrics, 168(9), 822-828

National Institute on Alcohol Abuse and Alcoholism. (2011). Alcohol Screening and Brief Intervention for Youth: A Practitioners Guide. NIH Publication No. 11-7805
 Russell M. (1994). New Assessment Tools for Risk Drinking During Pregnancy: T-ACE, TWEAK, and Others. Alcohol health and research world, 18(1), 55-61.

4. Yonkers KA, Gotman N, Kershaw T, Forray A, Howell HB, Rounsaville BJ. Screening for prenatal substance use: development of the Substance Use Risk Profile-Pregnancy scale. Obstet Gynecol. 2010 Oct;116(4):827-833.



Brief intervention/brief treatment

Brief intervention:

- Provide education for patients on risks of substance use.
- Motivate patients to reduce risky behavior.




Brief treatment

Involves setting goals for patient:

- Changing immediate behavior or thoughts about risky behavior
- Addressing longstanding problems with harmful drinking and drug misuse
- Helping patients with higher levels of disorder obtain more long-term care
- Brief treatment should generally accompany a referral to treatment



Brief treatment process



Referral to treatment

Referral is recommended when a patient meets the diagnostic criteria for substance use disorder, but diagnosing is not required for provider performing SBIRT:

- Patients are referred to a specialized treatment provider who can provide more long-term treatment for complex issues related to substance use.

Referrals may be made to several types of services (and more than one, if necessary):

- Outpatient counseling, individual, or group
- Acute treatment services (detox)
- Medication-assisted treatment
- Clinical stabilization services
- Support groups (AA, NA, Al-Anon)



Key resources

Substance Abuse and Mental Health Services Administration (SAMHSA)

<http://www.samhsa.gov/sbirt>

Centers for Medicare & Medicaid Services (CMS)

[SBIRT Under Medicare and Medicaid](#)



In closing

When applied correctly, SBIRT is very effective:

- Screening and brief interventions are both very effective for alcohol use.
- Screening is very effective for identifying illicit drug use.
- Referral to treatment should follow any positive screening for drug use.

SBIRT:

- Saves lives.
- Saves time.
- Saves money.





provider.wellpoint.com/dc

Coverage provided by Wellpoint District of Columbia, Inc.

DCWP-CD-058545-24 | May 2024